

## 2d: Treatment centres: Port Elizabeth

SANCA - PE

**Table 61: Proportion of treatment episodes (Port Elizabeth)**

	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%
SANCA PE	83	83	87	85	80	66	65	74	86
Welbedacht	11	8	9	10	7	15	12	11	9
Shepherd's Field	7	9	4	5	9	0	15	13	4
SANCA Thembilitsha	-	-	-	-	4	18	0	0	0
Bethesda	-	-	-	-	-	1	0	0	0
Family Outreach Ministries	-	-	-	-	-	-	8	2	<1
<b>Total no of persons treated</b>	<b>N=515</b>	<b>N=450</b>	<b>N=505</b>	<b>N=418</b>	<b>N=464</b>	<b>N=318</b>	<b>N=570</b>	<b>N=457</b>	<b>N=577</b>

A total of 577 patients were treated at the Port Elizabeth treatment centres.

**Table 62: First time admissions (Port Elizabeth)**

	Jul-Dec 2002	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%	%
<b>Yes</b>	81	74	67	76	70	78	78	74	74	79
<b>No</b>	19	26	33	24	30	22	22	26	26	21

79% of the treatment population reported that it was their first admission.

**Table 63: Types of treatment received (Port Elizabeth)**

	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%
<b>Inpatient</b>	18	18	15	18	20	36	26	26	17
<b>Outpatient</b>	82	82	85	82	80	64	74	74	83
<b>Both</b>	0	0	0	0	0	0	0	0	0

The majority of patients are seen on an outpatient basis (83%).

**Table 64: Referral sources (Port Elizabeth)**

	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%
Self/family/friends	39	42	45	45	42	48	49	50	50
Work/employer	17	18	16	18	21	20	20	16	16
Doctor/psychiatrist/nurse (health professional)	11	14	12	11	9	16	15	15	12
Religious body	2	2	1	2	1	<1	2	2	1
Hospital/clinic	10	7	9	9	10	3	6	5	5
Social services/welfare	8	6	4	7	7	3	3	4	4
Court/correctional services/police/lawyer	2	4	5	4	3	2	1	3	5
School	8	6	4	2	6	5	3	4	6
Other e.g. radio, Children's home, adverts	4	1	4	2	2	2	1	1	1

The majority of referrals are from family. Referrals from all sources have been fairly stable over the last 3 review periods. A slight decrease in referrals from health professionals was noted in this period.

**Table 65: Age distribution (Port Elizabeth)**

Years	Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%
Under 10	-	-	3	<1	0	0	1	<1	0	0	0	0
10-14	11	3	12	3	9	3	4	1	6	1	20	4
15-19	66	16	78	17	47	15	69	14	83	18	101	18
20-24	66	16	76	16	29	9	84	17	80	17	74	13
25-29	53	13	66	14	43	14	77	16	65	14	87	15
30-34	68	16	67	14	49	15	67	14	63	14	73	13
35-39	50	12	46	10	42	13	52	11	55	12	71	12
40-44	41	10	40	9	34	11	40	8	48	11	57	10
45-49	29	7	45	9	31	10	36	7	35	8	40	7
50-54	20	5	21	4	17	5	25	5	8	2	27	5
55-59	9	2	6	1	11	3	15	3	6	1	15	3
60-64	1	<1	7	1	2	<1	7	2	2	<1	5	1
65-69	3	<1	0	0	0	0	4	1	4	1	6	1
70-80	-	-	-	-	4	1	1	<1	-	-	0	0
Age unknown	1	<1	0	0	0	0	0	0	2	<1	1	<1

Almost a quarter of patients (22%) were younger than 20 years in the 1<sup>st</sup> half of 2007.

**Table 66: Population Profile (Port Elizabeth)**

	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%
<b>GENDER</b>									
Male	79	82	85	84	80	78	83	81	81
Female	21	18	15	16	20	22	17	19	19
<b>ETHNIC GROUP</b>									
Black	27	29	28	30	28	34	22	19	23
Asian	3	3	3	2	7	3	4	3	1
Coloured	46	42	41	43	44	42	41	46	46
White	24	26	27	25	25	21	33	32	30
<b>MARITAL STATUS</b>									
Married (living with spouse)	25	30	29	28	23	30	26	21	26
Living Together	6	6	8	9	5	12	10	8	6
Divorced	9	9	8	8	9	8	6	9	8
Widowed	1	2	2	2	2	3	2	1	1
Single	55	53	46	48	57	46	56	55	57
Separated/not living with spouse	-	-	-	-	-	<1	0	5	2
Other	-	-	-	2	1	<1	0	<1	0
<b>EMPLOYMENT STATUS</b>									
Working full-time	35	34	39	39	39	46	40	37	37
Working Part-time	3	5	5	3	3	4	4	4	6
Unemployed	33	40	33	35	33	26	36	39	33
Intern/Apprentice	<1	3	<1	<1	0	<1	1	1	0
Learner/Student	22	15	14	15	17	16	14	14	19
Disabled/Boarded	2	1	2	2	1	1	2	1	2
Housewife	2	1	1	<1	2	2	1	1	<1
Pension/Retired	2	2	2	2	2	3	1	2	3
Other	-	2	0	2	3	<1	1	1	<1

Table 66 shows the population profile of patients attending treatment centres in Port Elizabeth. The single adult male remains the dominant profile of the individual in treatment. The majority of patients are unemployed (33%) or students (19%).

**Table 67: Primary substance of abuse (Port Elizabeth)**

	Jul-Dec 2002	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%	%
Alcohol	55	46	51	48	45	47	59	46	39	48
Dagga	13	16	12	15	13	12	14	16	15	17
Dagga/Mandrax	25	32	26	24	25	20	9	9	9	10
Cocaine	<1*	2*	2*	6*	6	3	3	22*	4	16*
Crack					3	9	9		20	
Ecstasy	1	<1	2	3	1	<1	0	2	1	0
OTC/PRE <sup>1</sup>	4	5	5	3	3	5	6	3	4	5
Polysubstance	0	<1	<1	0	0	0	0	0	0	0
Heroin	<1	0	0	2	3	2	<1	2	3	1
Ephedrine	-	-	1	0	0	0	0	0	0	0
Inhalants	-	-	1	0	0	<1	0	<1	<1	<1
Nicotine	0	0	<1	0	0	0	0	0	<1	0
Methamphetamine ('Tik')	-	-	-	-	-	1	0	1	4	2
Khat	-	-	-	-	-	<1	0	0	0	<1
Other	0	<1	0	0	<1	0	0	0	0	<1

\*Crack/cocaine combined

<sup>1</sup>Over the counter or prescription medicines

Alcohol remains the primary substance of abuse (48%), followed by dagga and cocaine.

**Table 68: Frequency of use of primary substance of abuse (Port Elizabeth)**

	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%
Daily	52	59	58	62	60	48	55
2-6 days per week	37	33	32	31	27	35	30
Once a week or less	8	8	9	5	7	9	13
Not used in past month	2	<1	1	1	6	8	2

**Table 69: Age at which primary substance of abuse first used or experimented with (Port Elizabeth)**

Years	Jan-Jun 2004		Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
5-10	-	-	3	<1	11	2	4	1	1	<1	6	1	-	-
11-19	339	67	275	66	290	63	178	57	280	58	267	58	348	60
20-29	126	25	103	25	117	25	85	27	151	31	126	28	150	26
30-39	26	5	26	6	39	8	26	8	39	8	40	9	58	10
40-49	11	2	7	3	13	3	8	3	8	2	13	3	13	2
50-59	3	<1	1	<1	2	<1	5	2	2	<1	3	<1	5	<1
60-69	-	-	-	-	-	-	-	-	-	-	-	-	3	<1
Age unknown	-	-	3	<1	-	-	7	2	1	<1	2	<1	-	-

The majority of users (primary substance of abuse) reported having started abusing drugs between the ages of 10-19 years.

**Table 70: Gender by primary substance of abuse (Port Elizabeth)**

	%													
	Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007			
	M	F	M	F	M	F	M	F	M	F	M	F		
Alcohol	84	16	75	25	77	23	81	19	78	22	81	19		
Dagga/Mandrax	94	6	95	5	97	3	95	5	95	5	93	7		
Dagga	83	17	93	7	93	7	96	4	87	13	92	8		
Crack	100	0	91	9	82	18	78	22	85	15	77 <sup>+</sup>	23 <sup>+</sup>		
Cocaine	80	20	83	17	75	25	76	24	72	28				
Ecstasy	67	33	100	0	0	0	86	14	100	0	0	0		
OTC	21*	79*	60	40	20	80	67	33	50	50	46*	54*		
PRE			12	88	14	86	33	67	38	62				
Heroin	75	25	68	33	100	0	100	0	54	46	50	50		
Inhalants	0	0	0	0	0	0	100	0	100	0	67	33		
Methamphetamine ('Tik')	-	-	50	50	0	0	80	20	89	11	30	70		
Other	-	-	-	-	-	-	-	-	-	-	100	0		
Khat	-	-	-	-	-	-	-	-	-	-	100	0		

\*OTC/PRE combined    <sup>+</sup> Crack/cocaine combined

Very few Mandrax patients are female, whereas the majority of patients treated for over-the-counter or prescription medication abuse are females.

**Table 71: Race by primary substance of abuse (Port Elizabeth)**

	BLACK			COLOURED			ASIAN			WHITE		
	%			%			%			%		
	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
Alcohol	31	29	30	38	43	38	2	<1	1	29	27	30
Dagga/ Mandrax	24	28	37	69	61	58	0	2	0	7	9	5
Dagga	23	23	21	46	49	62	4	3	1	27	26	16
Crack	4	4	5 <sup>+</sup>	45	54	43 <sup>+</sup>	9	7	2 <sup>+</sup>	42	34	50 <sup>+</sup>
Cocaine	0	0		12	17		0	0		88	83	
Ecstasy	0	0	0	29	20	0	0	0	0	71	80	0
OTC	0	0	0*	0	0	54*	0	0	0*	100	100	46*
PRE	17	6		58	31		8	13		17	51	
Heroin	12	0	0	13	15	0	0	8	0	75	77	100
Metham- phetamine (‘Tik’)	0	0	0	20	61	60	40	0	0	40	39	40
Khat	0	0	0	0	0	0	0	0	0	0	0	100
Inhalants	100	0	33	0	0	67	0	0	0	0	100	0
Other	0	0	0	0	0	100	0	0	0	0	0	0

\*OTC/PRE combined <sup>+</sup>Crack/cocaine combined

There was an almost equal distribution of Black, Coloured and White alcohol patients in the 1<sup>st</sup> half of 2007, while cocaine patients were mostly White or Coloured.

**Table 72: Average/Mean age by Primary Substance (Port Elizabeth)**

	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	Mean age of Patients	Mean age of Patients	Mean age of Patients	Mean age of Patients	Mean age of Patients	Mean age of Patients
Alcohol	38	36	38	39	37	37
Dagga/ Mandrax	26	25	25	26	31	27
Dagga	21	22	21	21	20	20
Ecstasy	-	-	-	24	28	-
Cocaine	27	28	23	29	28	28 <sup>+</sup>
Crack	25	27	27	26	27	
OTC	33*	39*	41	35	41	41*
PRE			40	37	39	
Heroin	25	28	24	25	25	29
Inhalants	-	-	-	49	14	-
Methamphetamine (‘Tik’)	-	24	-	29	22	26

\*OTC/PRE Combined <sup>+</sup>Cocaine/Crack combined

Patients treated for alcohol, prescription and over-the-counter medicines are on average older than those treated for other more commonly used drugs such as cannabis, crack and Mandrax.

**Table 73: Mode of use for primary drug (Port Elizabeth)**

	Jan-Jun 2004		Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
<b>Swallowed</b>	276	55	215	51	239	52	206	65	241	50	206	45	300	52
<b>Smoked</b>	215	43	176	42	204	44	111	35	218	45	232	51	258	45
<b>Snorted</b>	9	2	20*	5*	14*	3*	0	0	20*	4*	13*	3*	13*	2*
<b>Sniffed</b>	0	0					0	0						
<b>Injected</b>	5	1	7	2	7	2	1	<1	3	1	5	1	4	<1
<b>Other/Combination</b>	-	-	-	-	-	-	-	-	-	-	1	<1	2 <sup>+</sup>	<1

\*Snorted/sniffed combined    <sup>+</sup>Smoke/inject combined

Swallowing and smoking remain the more common methods of using drugs.

**Table 74: Secondary substance of abuse (Port Elizabeth)**

	Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%
<b>Alcohol</b>	46	34	52	31	29	36	45	25	39	21	46	28
<b>Dagga/Mandrax</b>	17	13	37	22	6	7	28	16	32	17	20	12
<b>Dagga</b>	23	17	22	13	21	26	36	20	47	26	35	21
<b>Cocaine</b>	5	4	3	2	2	3	19	11	16	9	38*	23*
<b>Crack</b>	17	13	23	14	14	18	19	11	23	12		
<b>Ecstasy</b>	13	10	6	4	1	1	10	6	10	5	8	5
<b>LSD</b>	0	0	0	0	1	1	4	2	3	2	0	0
<b>OTC</b>	4	3	1	<1	2	3	0	0	2	1	7 <sup>+</sup>	4 <sup>+</sup>
<b>PRE</b>	2	2	9	5	4	5	0	0	3	2		
<b>Heroin</b>	5	4	1	<1	0	0	7	4	4	2	8	5
<b>Inhalants</b>	0	0	12	7	0	0	0	0	0	0	1	<1
<b>Khat</b>	1	1	0	0	0	0	6	3	1	<1	0	0
<b>Polysubstance</b>	0	0	1	<1	0	0	0	0	0	0	0	0
<b>Methamphetamine ('Tik')</b>	-	-	-	-	-	-	-	-	4	2	3	2

\*Cocaine/Crack combined    <sup>+</sup> OTC/PRE Combined

**Table 75: Frequency of use of secondary substance of abuse (Port Elizabeth)**

	Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%
Daily	27	20	44	27	29	36	60	34	45	25	39	24
2-6 times a week	52	39	75	45	34	43	51	29	62	35	62	38
Once a week or less frequent	41	31	38	23	13	16	41	23	39	21	44	27
Not used in past month	13	10	9	5	4	5	28	14	34	19	20	12

24% of patients reported to using their secondary substance of abuse daily.

**Table 76: Source of payment (Port Elizabeth)**

	Jul-Dec 2002	Jan-Jun 2003	Jul-Dec 2003	Jan-Jun 2004	Jul-Dec 2004	Jan-Jun 2005	Jul-Dec 2005	Jan-Jun 2006	Jul-Dec 2006	Jan-Jun 2007
	%	%	%	%	%	%	%	%	%	%
<b>Self</b>	35	28	34	40	40	37	37	41	37	37
<b>Medical Aid</b>	24	18	10	21	19	15	20	12	10	9
<b>Family</b>	30	31	35	27	26	37	32	36	38	40
<b>Friends</b>	1	1	<1	<1	<1	<1	1	<1	<1	<1
<b>Employer</b>	2	4	5	4	5	5	7	7	9	9
<b>Unknown</b>	8	11	14	5	5	3	4	2	4	3
<b>Other</b>	<1	6	0	4	4	3	<1	1	2	2
<b>State</b>	-	-	<1	<1	<1	<1	<1	0	<1	0

Family and 'self' are the most common sources of payment.

**Table 77: Treatment population – suburb of residence (Port Elizabeth)**

	Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%
<b><i>METRO SUBSTRUCTURE</i></b>												
Bethelsdorp, Gelvandale, Korsten, Malabar	127	30	144	31	94	39	123	26	171	37	158	27
Roadhouse, New Brighton, Motherwell	92	22	77	16	52	16	65	13	52	11	118	20
Uitenhage, Despatch	74	18	69	15	38	12	49	10	54	12	102	18
North End, Mount Road, Central	42	10	41	9	48	15	25	5	19	4	46	8
Walmer, Kabega Park, Summerstrand	41	10	80	17	22	7	81	17	40	9	51	9
Port Elizabeth (other parts)	-	-	-	-	-	-	80	17	66	14	45	8
<b><i>OTHER PARTS OF THE PROVINCE</i></b>												
George, King Williams Town, Humansdorp, Maclear and East London	24	6	46	10	64	20	54	11	53	12	11	2
<b>Unknown</b>	18	4	1	<1	0	0	5	1	2	<1	25	4
<b>Other Provinces</b>	-	-	-	-	-	-	-	-	-	-	21	4

**Table 78: Tertiary substance of abuse (Port Elizabeth)**

	Jan-Jun 2004		Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Alcohol	10	22	9	23	9	25	2	15	18	24	16	22	15	36
'White Pipe'	7	16	7	18	6	15	1	8	6	8	10	13	1	2
Dagga	4	9	4	11	2	6	1	8	4	5	12	16	9	21
Cocaine	6	13	2	5	4	11	3	23	8	11	6	8	13	31
Crack	8	18	4	11	4	11	1	8	9	12	9	12		
Heroin	1	2	2	5	1	2	1	8	9	12	0	0	0	0
Ecstasy	6	13	8	21	8	22	4	31	9	12	9	12	4	10
Ecstasy/ Dagga	-	-	-	-	1	3	0	0	0	0	0	0	0	0
OTC/PRE	2	4	1	3	1	3	0	0	3	4	4	6	0	0
Metham- phetamine (‘Tik’)	-	-	-	-	-	-	-	-	5	7	3	4	0	0
Moonflower	-	-	1	3	0	0	0	0	0	0	0	0	0	0
LSD	1	2	0	0	0	0	0	0	3	4	5	7	0	0
Other	-	-	1	3	0	0	0	0	0	0	0	0	0	0

## ***SUBSTANCE ABUSE FOR PERSONS UNDER 20 YEARS***

**Table 79: Primary substance of abuse of patients younger than 20 years (Port Elizabeth)**

	Jan-Jun 2004		Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Alcohol	9	10	8	10	24	27	13	23	17	23	14	16	32	26
Dagga/ Mandrax	32	37	30	39	27	30	9	16	8	11	4	5	11	9
Dagga	37	42	32	41	31	35	29	52	35	47	47	53	63	52
Cocaine	1	1	4	5	1	1	1	2	0	0	2	2	9 <sup>+</sup>	7 <sup>+</sup>
Crack	1	1	2	3	4	4	4	7	9	12	14	16		
Ecstasy	5	6	1	1	0	0	0	0	2	3	0	0	0	0
Heroin	1	1	0	0	0	0	0	0	1	1	0	0	1	<1
PRE	1	1	1	1	0	0	0	0	1	1	0	0	0	0
Inhalants	0	0	0	0	1	1	0	0	0	0	1	1	3	3
Metham- phetamine (‘Tik’)	-	-	-	-	-	-	-	-	1	1	6	7	1	<1
Other	-	-	-	-	-	-	-	-	-	-	-	-	1	<1
<b>TOTAL</b>	<b>87</b>	<b>100</b>	<b>78</b>	<b>100</b>	<b>90</b>	<b>100</b>	<b>56</b>	<b>100</b>	<b>74</b>	<b>100</b>	<b>89</b>	<b>100</b>	<b>121</b>	<b>100</b>

<sup>+</sup>Cocaine and Crack combined

Dagga remains the most common primary substance for patients younger than 20 years, followed by alcohol and crack.

**Table 80: Gender of patients abusing substances younger than 20 years (Port Elizabeth)**

	Jan-Jun 2004		Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
<b>Male</b>	73	84	68	87	67	75	45	80	63	85	69	78	97	80
<b>Female</b>	14	16	10	13	23	25	11	20	11	15	20	22	24	20

**Table 81: Race of patients younger than 20 years (Port Elizabeth)**

	Jan-Jun 2004		Jul-Dec 2004		Jan-Jun 2005		Jul-Dec 2005		Jan-Jun 2006		Jul-Dec 2006		Jan-Jun 2007	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
<b>Black</b>	28	32	32	41	28	31	24	43	16	22	13	15	30	25
<b>Asian</b>	2	3	3	4	3	3	0	0	1	1	3	3	2	2
<b>Coloured</b>	35	40	29	37	52	58	27	48	43	58	47	52	73	60
<b>White</b>	22	25	14	18	7	8	5	9	14	19	26	30	16	13
<b>TOTAL</b>	<b>87</b>	<b>100</b>	<b>78</b>	<b>100</b>	<b>90</b>	<b>100</b>	<b>56</b>	<b>100</b>	<b>74</b>	<b>100</b>	<b>89</b>	<b>100</b>	<b>121</b>	<b>100</b>

The majority of youths seeking treatment are from the Coloured population.